This little piggy coughed all the way home

Dr. J. Lowe[1]

[1]University of Illinois Urbana Champaign

The threat of pandemic influenza from swine-origin virus presents a complex epidemiological challenge that requires comprehensive understanding of both viral movement within pig populations and the pathways through which these viruses transmit to humans. This presentation examines the intricate network structures that facilitate influenza A virus (IAV) transmission between swine and human populations.

The North American swine industry operates through two distinct but interconnected ecosystems that create different transmission risk profiles. The exhibition swine ecosystem, centered around agricultural fairs and show pigs, involves small seasonal herds with high levels of human-pig contact, particularly among children and fair participants. These animals are frequently mixed at fairs throughout their lives, creating opportunities for viral reassortment and amplification. In contrast, the commercial swine ecosystem consists of large year-round operations with professional management, limited human contact, and cohorts of pigs that remain stable until harvest.

Network analysis reveals that while these ecosystems appear separate, they are connected through proximity and shared human resources. The presentation demonstrates seasonal patterns of transmission risk, with show pig networks showing increased connectivity during fair season from May through September. During this period, widespread mixing of animals from multiple small breeding operations creates ideal conditions for viral evolution and cross-species transmission.

Historical evidence indicates that all current pig influenza A viruses originated from human populations, while pigs subsequently serve as a source of IAV for humans with potentially less genetic diversity than human-to-pig transmission. This bidirectional flow creates opportunities for viral reassortment and the emergence of novel strains with pandemic potential.

There are critical knowledge gaps that limit effective pandemic preparedness: insufficient understanding of commercial network interconnections, incomplete mapping of show pig networks, and limited comprehension of how these two ecosystems interface. The multidimensional nature of these connections—involving animals, people, fomites, trucks, and wildlife—presents measurement and modeling challenges that extend beyond traditional single-host pathogen surveillance systems.

Proposed intervention strategies focus on vaccination and testing protocols for both pig populations and fair attendees, though implementation requires better understanding of viral strain diversity and circulation patterns. Effective pandemic prevention requires interdisciplinary collaboration, acknowledging that "the pig guys are not going to solve this alone."

This work underscores the urgent need for comprehensive network mapping, enhanced surveillance systems, and coordinated intervention strategies to minimize the risk of swine-origin influenza pandemics in human populations.